arXiv:math/0511365v4 [math.CO] 15 Aug 2006

ON SQUENCE OF QUADRATIC EXPONENTIAL FUNCTIONS CONNECTED TO $\mathbb{N}$

Arni S.R. Srinivasa Rao
Mathematical Institute, University of Oxford,
24-29 St Giles', Oxford, OX1 3LB, England.
Tel: +44-1865-283886
Fax: +44-1865-270515
email: arni@maths.ox.ac.uk

ABSTRACT. Link between quadratic exponential functions and natural numbers is investigated through classical analysis arguments. An useful mid-point theorem is proved.

Key words: natural numbers mapping, mean value theorem.
MSC: 26A09,26E99

1. BACKGROUND AND MOTIVATION

A correspondence between natural numbers and sequence of quadratic exponential functions is investigated in an elementary way. Never before the properties of these functions were analyzed in this fashion. Let f be a quadratic exponential function with (domain) $D(f) = \mathbb{N}$. For a given combination (say $\mathcal{C}_1$) of integer parameters of f, we establish here that the mapping of $1 \in D(f)$ will be same as mapping of some integer $n \in D(f)$ (for the same combination of parameters). If we change the combination of parameters (say $\mathcal{C}_2$), then the resultant mapping of 1 will be same as mapping of some $n_1 \in D(f)$. Here $n \neq n_1$ and $f(1)$ for $\mathcal{C}_1$ is not equal to $f(1)$ for $\mathcal{C}_2$ i.e. $f_{\mathcal{C}_1}(1) \neq f_{\mathcal{C}_2}(1)$. See figure 1.1. We construct such quadratic exponential sequence of numbers and try to link them to the natural numbers. The readers will also see that for some $n_k \in D(f)$, the distance from 1 to n_k

will be equal to $f(n_k)$ in certain conditions. We apply these facts to establish further interesting properties of convergence and derivative of f. Using Rolle's theorem we show that $f'(\theta) = 0$ for the mid-point $\theta \in (1, n_k)$.

Consider the quadratic function, $Q_1(x) = u_1x^2 + v_1x + w_1$ where $u_1 > 0, v_1, w_1 \in \mathbb{R}$. Suppose $u_1 = \frac{m_1}{2}, v_1 = \frac{n_1}{2}$ and $m_1(>0), n_1, c_1 \in \mathbb{N}$, $m_1 + n_1$ is even, then $Q_1(x)$ is an integer for an integer x [1, 2]. The versatile features of quadratic function when its coefficients are positive integers or real numbers have been popular in modeling natural sciences [3]. When quadratic function is taken as an exponent to the irrational number e, then the resultant function is called quadratic exponential function. Functions from such family were widely established tools in modeling biological data [4, 5]. These functions can even mimic properties of Gaussian probability functions [5]. Suppose $f(x) = \exp\left(ux^2 + vx\right)$, $v = (-1)^{2k-1}mu^{2k-1}$, $u, m, k \in \mathbb{N}$, then for $A = \frac{u+v}{u}$, it was proved that $f(1) = f(|A|)$ [6]. In fact this statement was also proved there for $k = 1$ and $u \in \mathbb{R} - \{0\}$, $m \in \mathbb{N}$. We use these results and establish few interesting properties of such exponential function. By using Rolle's theorem we show that the derivative will be zero at the mid-point of the interval $[1, |A|]$.

Note that obviously $Q(1) = Q(|A|)$ when $Q(x) = ux^2 + vx$. Also note $|A| = 1 - mu^{2k-2}$ if $(1 \geq mu^{2k-2})$ or $|A| = mu^{2k-2} - 1$ if $(1 < mu^{2k-2})$. Try and see that this absolute value function is necessary for deriving the results in this work. We begin with some simple results.

Lemma 1.1. *$|A|$ is always a positive integer for any combination of $\{k, m, u\} \in \mathbb{N}$.*

Proof. $|A| = \left|\frac{u+(-1)^{2k-1}mu^{2k-1}}{u}\right| = mu^{2k-2} - 1$. $u \in \mathbb{N} \Rightarrow u^{2k-2} \in \mathbb{N}$ (since $m, u \in \mathbb{N}$. Therefore $|A| \in \mathbb{N}$. $\square$

Additionally, whenever $u = \left(\frac{n+1}{m}\right)^{\frac{1}{2k-2}}$, $n \in \mathbb{N}$, then $u^{2k-2} = \frac{n+1}{m}$, which means $\left|\frac{u+v}{u}\right| = n$. For that matter $|A|$ could be equal to every natural number for a suitable combination of $\{k, m, u\} \in \mathbb{N}$. For example, $\{k = 1, m = 2, u = 2\}$ gives $|A| = 1$, $\{k = 1, m = 3, u = 2\}$ gives $|A| = 2$, $\{k = 1, m = 4, u = 2\}$ gives $|A| = 3$, $\{k = 3, m = 10, u = 8\}$ gives $|A| = 40959$. We know that



FIGURE 1.1. $f_p(1) = f_p(n_k)$ for the p^{th} combination of parameters of f and $n_k \in \mathbb{N}$.

$f(1) = f(|A|)$. Therefore, we modify the previous result and state that as follows:

Lemma 1.2. *For any combination of* $\{k,m,u\} \in \mathbb{N}$, *there corresponds a* $f(1)$ *such that* $f(1) = f(n_k)$ *for some* $n_k \in \mathbb{N}$.

Proof. From lemma 1.1, for every $|A|$ there corresponds a $n \in \mathbb{N}$ and $f(1) = e^{u\left(1-mu^{2k-2}\right)} = f(|A|)$. Thus $f(1) = f(n_k)$ for some $n_k \in \mathbb{N}$. □

Note 1.3. The domain of f is $\mathbb{N}$. The value of $f(1)$ is not same for every combination of $\{k,m,u\} \in \mathbb{N}$. Readers are suggested to keep this in mind for understanding the results presented in this work.

2. LINKING NATURAL NUMBERS AND EXPONENTIAL FUNCTION

Theorem 2.1. *When N is even,* $v = (-1)^{Nk-1} mu^{Nk-1}$ *and* $\{m,u,k,N\} \in \mathbb{N}$, *then it follows that* $f(1) = f(n_k)$, *whenever* $|A| \in \mathbb{N}$ *and for* $n_k \in \mathbb{N}$.

Proof. $f(1)=e^{u\left(1-mu^{Nk-2}\right)}$ and $|A|=\left|\frac{u+(-1)^{Nk-1}mu^{Nk-1}}{u}\right|=mu^{Nk-2}-1$. Therefore, $|A|\in N$ for $\{m,u,k,N\}\in\mathbb{N}$. Now,

$$\begin{aligned} f(|A|) &= e^{u\left(mu^{Nk-2}-1\right)^2+v\left(mu^{Nk-2}-1\right)} \\ &= e^{u\left(m^2u^{2Nk-4}+1-2mu^{Nk-2}\right)-m^2u^{2Nk-3}+mu^{Nk-1}} \\ &= e^{u\left(1-mu^{Nk-2}\right)} \end{aligned}$$

Therefore, $f(1)=f(|A|)$. Since $|A|$ consists of every element of $\mathbb{N}$, it follows that $f(1)=f(n)$. □

Theorem 2.2. *When N is odd, k is even,* $v=(-1)^{Nk-1}mu^{Nk-1}$ *and* $\{m,u,k,N\}\in\mathbb{N}$*, then it follows that* $f(1)=f(n_k)$*, whenever* $|A|\in\mathbb{N}$ *and for* $n_k\in\mathbb{N}$*.*

Proof. When N is odd and k is even $|A|=\left|\frac{u+(-1)^{Nk-1}mu^{Nk-1}}{u}\right|=mu^{Nk-2}-1$. The rest of the proof can be deduced from theorem 2.1. □

Remark 2.3. Suppose lemma 1.1 holds and f_p denotes p^{th} combination of $\{m,u,k\}$, say $\left\{m_p,u_p,k_p\right\}$, then

$$\begin{aligned} f_1(1) &= f_1(2)\neq f_1(3)\neq\ldots\neq f_1(n)\neq\ldots \\ f_2(1) &\neq f_2(2)=f_2(3)\neq\ldots\neq f_2(n)\neq\ldots \\ &\vdots \\ f_p(1) &= f_p(2)\neq f_p(3)\neq\ldots\neq f_p(n)=f_p(n+1)\neq\ldots \\ &\vdots \end{aligned}$$

Remark 2.4. Given the theorem 2.1, the distance $\mathcal{D}$, from $|A|$ to 1 is equal to $f(|A|)$ if $m=\mathcal{Z}(1-\mathcal{Z})^{Nk-2}/\{log(\mathcal{Z}-2)\}^{Nk-2}$. Here $\mathcal{Z}=mu^{Nk-2}$.

Remark 2.5. Suppose $\mathcal{D}_p$ be the logarithamic distance from 1 to $|A|$ for the p^{th} combination of parameters, then

$$\begin{aligned}
\mathcal{D}_p &= \log\{\mathcal{Z}_p - 2\} \\
&= \log\left\{\mathcal{Z}_p\left(1-\frac{2}{\mathcal{Z}_p}\right)\right\} = \log\mathcal{Z}_p + \log\left(1-\frac{2}{\mathcal{Z}_p}\right)
\end{aligned}$$

$\mathcal{D}_p$ converges for $\mathcal{Z}_p > 2$.

Remark 2.6. $f_m(1) = f_m(n)$ is unique for each $n \in \mathbb{N}$ and $\{m_p, u_p, k_p\} \in \mathbb{N}$.

Remark 2.7. Suppose N_{p+i} is even for $\begin{cases} p = 1,2,3,\dots \\ i = 0,1,2,\dots \end{cases}$. If $|A_p| > |A_{p-1}|$ then $f_p(1) < f_{p-1}(1)$ for all $p = 2,3,4,\dots$. This is also explained through figure 3.1.

Remark 2.8. The number of pairs $\{f_m(1),\ f_m(n)\}$ that satisfy remark 2.6 are countable.

Theorem 2.9. *When N_{p+i} is even for $\begin{cases} p = 1,2,3,\dots \\ i = 0,1,2,\dots \end{cases}$, if $|A_{p+1}| > |A_p|$ then the sequence $\{f_p(1)\}_{p=1,2,3,\dots} \in \mathcal{B}_\sigma(0)$ for $p > M \in \mathbb{N}$. Here $\mathcal{B}_\sigma(0) = \{b \in \mathbb{R}^+ : |b-0| < \sigma\}$ for $\sigma > 0$.*

Proof. $|A_{p+1}| > |A_p| \Rightarrow m_{p+1}u^{N_{p+1}k_{p+1}-2} - 1 > m_p u^{N_p k_p - 2} - 1$.

$$\begin{aligned}
\Rightarrow \left(1 - m_{p+1}u^{N_{p+1}k_{p+1}-2}\right) &< \left(1 - m_p u^{N_p k_p - 2}\right) \\
\Rightarrow u_{p+1}\left(1 - m_{p+1}u^{N_{p+1}k_{p+1}-2}\right) &< u_p\left(1 - m_p u^{N_p k_p - 2}\right) \\
\Rightarrow e^{u_{p+1}\left(1 - m_{p+1}u^{N_{p+1}k_{p+1}-2}\right)} &< e^{u_p\left(1 - m_p u^{N_p k_p - 2}\right)}
\end{aligned}$$

This implies $f_{p+1}(1) < f_p(1)$. We know that $e^{-p} \to 0$ as $p \to \infty$. Thus $f_p(1) \in \mathcal{B}_\sigma(0)$. □

Corollary 2.10. *Since $f_p(1) \in \mathcal{B}_\sigma(0)$, it follows from remark 2.3 that $\{f_1(2), f_2(3), \dots, f_p(n)\ , f_{p+1}(n+1), \dots\}$ is a convergent sequence.*

3. MID-POINT THEOREM

Theorem 3.1. *(Mid-point theorem) It is easy to verify that f is continuous on $[1, |A|]$ and differentiable on $(1, |A|)$, and from lemma 1.2 we have $f(1) =$*

$f(|A|)$*, so by Rolle's theorem there exists a* $\theta \in (1,|A|)$ *such that* $f'(\theta) = 0$*. Then this* θ *is the mid-point of the interval* $I = [1,|A|]$*.*

Proof. We have $|A| = mu^{2k-2} - 1$. Mid-point of the interval I is $\frac{m}{2}u^{2k-2}$.

$f'(\theta) = e^{u\theta^2+v\theta}(2u\theta+v) = f(\theta)(2u\theta+v)$. Since $f'(\theta) = 0$, this means obviously $2u\theta+v = 0$, because $f(\theta) \neq 0$. Therefore $\theta = \frac{-v}{2u} = \frac{m}{2}u^{2k-2}$. Hence θ is the mid-point of the interval I. Fors a numerical example, consider $\{k = 3,\ m = 10, u = 8\}$ as in section 1. For this combination $|A| = 40959$, mid-point of the interval is 20480 and $f'(20480) = 0$. ☐

Theorem 3.2. *Suppose* $\delta_1 < \delta_2 < \ldots\delta_n$*, where* $\delta_i(> 0) \in I$*. Then for a given combination of* $\{k,m,u\}$*,* $f(1+\delta_1) > f(1+\delta_2) > \ldots f(1+\delta_n) = f(|A|-\delta_n) < \ldots < f(|A|-\delta_2) < f(|A|-\delta_1)$ *if and only if* $(1+\delta_n)$ *is a mid-point of* I*, where* $\delta_n = \frac{m}{2}u^{2k-2} - 1$*.*

Proof. Verify easily that $f(1+\delta) = f(|A|-\delta)$ for $\delta(> 0) \in I$. Consider

$$f(1+\delta_1) > \ldots > f(1+\delta_n) \quad = \quad f(|A|-\delta_n) < \ldots < f(|A|-\delta_1) \tag{3.1}$$

By theorem 3.1 we know $f'(\theta) = 0$ for $\theta \in I$. Hence $(1+\delta_n)$ is a mid-point.

To prove converse we begin as follows. Since mid-point of the interval I is $\frac{m}{2}u^{2k-2}$, we have

$$\begin{aligned} f(\frac{m}{2}u^{2k-2}) &= e^{4(\frac{m}{2}u^{2k-2})^2 - mu^{2k-1}\frac{m}{2}u^{2k-2}} \\ &= e^{-\frac{m^2}{4}u^{4k-3}} \end{aligned} \tag{3.2}$$

Now for given $\delta_n = \frac{m}{2}u^{2k-2} - 1$, we can verify that

$$f(1+\delta_n) = f(|A|-\delta_n) \quad = \quad e^{-\frac{m^2}{4}u^{4k-3}} \tag{3.3}$$

Since $\delta_1 < \delta_2 < \ldots\delta_n$ and by equations 3.2 and 3.3, the result 3.1 is straightforward. ☐

For large value of the distance function $\mathcal{D}$ defined in remark 2.4, the shape of f look like the alphabet U [6]. Suppose instead of positive integer,



FIGURE 3.1. Relation between $\left|A_p\right|$ and $f_p(1)$. Vertical lines corresponding to $\left|A_p\right|$ are lengths of $f_p(1)$ for each p.

let $u \in \mathbb{Z}^-$and other parameters k, m remain as before, and if we denote resulting function as g, then the shape of g was shown to have mirror image of U [6]. Based on this information and from theorem 3.2, we state the following corollary.

Corollary 3.3. *Suppose* $\delta_1 < \delta_2 < \ldots\delta_n$, *where* $\delta_i(>0) \in I$. *Then for a given combination of* $\{k,m,u\}$, $g(1+\delta_1) < g(1+\delta_2) < \ldots g(1+\delta_n) = g(|A|-\delta_n) > \ldots > g(|A|-\delta_2) > g(|A|-\delta_1)$ *if and only if* $(1+\delta_n)$ *is a mid-point of* I, *where* $\delta_n = \frac{m}{2}u^{2k-2} - 1$.

Readers can also verify Darboux's theorem for f on the intervals $[1,\theta]$ and $[\theta,|A|]$ for some $f'(1) > \beta_1 > f'(\theta)$ or $f'(\theta) < \beta_2 < f'(|A|)$ such that $f'(\beta_1) = \alpha_1$ or $f'(\beta_2) = \alpha_2$ for $\alpha_1 \in [1,\theta]$ and $\alpha_2 \in [\theta,|A|]$.

In general, results on dynamics and periodic properties for the quadratic function of the form $x^2 + K$ [7] and periodic properties of natural numbers [8] can be found. However, this present note is basically deals with a correspondence between natural numbers, quadratic exponential function, and the convergence of such functions mapped on natural numbers constructed using $|A|$.

REFERENCES

[1] Pall, G. On sums of two or four values of a quadratic function of x. Trans. Amer. Math. Soc. 34 (1932), no. 1, 98–125.

[2] Pall, G. The structure of the number of representations function in a binary quadratic form. Trans. Amer. Math. Soc. 35 (1933), no. 2, 491–509.

[3] Ojha, V. P.; Pandey, H. A population growth model with the marriage rate as a quadratic function of time. J. Nat. Acad. Math. India 7 (1989), no. 2, 99–104.

[4] Cox, D.R and Wermuth, N. A note on the quadratic exponential binary distribution. Biometrika 81 (1994), no. 2, 403–408.

[5] McCullagh, P. Exponential mixtures and quadratic exponential families. Biometrika 81 (1994), no. 4, 721–729.

[6] Rao, A.S.R.S. "U" type functions. Bull. Inform. Cybernet. 35 (2003), no. 1-2, pp 35–39.

[7] Walde, R and Russo, P. Rational periodic points of the quadratic function $Q_c(x) = x^2 + c$. Amer. Math. Monthly 101 (1994), no. 4, pp318–331.

[8] Fine, N. J. Classes of periodic sequences. Illinois J. Math. 2 1958 285–302.